arXiv:1211.0874v2 [hep-ph] 13 Nov 2012

Transition Form Factors and Mixing of Pseudoscalar Mesons from Anomaly Sum Rule

Yaroslav Klopot,[1, *] Armen Oganesian,[1, 2, †] and Oleg Teryaev[1, ‡]
[1]*Joint Institute for Nuclear Research, Dubna 141980, Russia*
[2]*Institute of Theoretical and Experimental Physics, Moscow 117218, Russia*

The transition form factors of π^0, η, η' mesons are investigated by means of the anomaly sum rule – an exact nonperturbative relation which follows from the dispersive representation of the axial anomaly. Considering the problem of contributions of operators originated from a non-(local) OPE we found that they are required by the available set of experimental data, including the most recent data from the BELLE collaboration (which, if taken alone, can be described without such contributions, although do not exclude them). In this approach we analyzed the experimental data on η and η' meson transition form factors and obtained the constraints on the decay constants and mixing parameters.

We dedicate this paper to the memory of our friend and colleague Aleksander Bakulev.

PACS numbers: 11.55.Fv, 11.55.Hx, 14.40.Be, 13.60.Le

I. INTRODUCTION

The phenomenon of axial anomaly [1, 2] plays an important role in non-perturbative QCD and hadronic physics. The axial anomaly is known to govern the two-photon decays of the π^0, η and η' mesons and is usually considered for a case of real photons. However, the dispersive form of it [3] can be considered for virtual photons also [4–6] leading to a number of interesting applications.

One of the consequences of the dispersive approach to axial anomaly is a so-called anomaly sum rule (ASR) [5]. It gives, in particular, a complementary way to describe the π^0 [7] and the η, η' [8, 9] transition form factors (later developed also in [10, 11]) at all Q^2 even beyond the QCD factorization. This is especially important in view of the recent experimental studies of the $\gamma\gamma^* \to \pi^0(\eta, \eta')$ transitions [12–14]. In particular, the pion transition form factor, measured by the BABAR collaboration [12], revealed unexpectedly large values in the range of $Q^2 = 10 - 35$ GeV^2 resulting in an excess of the pQCD predicted limit [15] $Q^2 F_{\pi\gamma} \to \sqrt{2} f_\pi$, $f_\pi = 0.1307$ GeV. This striking result attracted a lot of interest and motivated extensive theoretical investigations. As a result, the transition form factors were (re)investigated using the framework of light cone sum rules [16–20], including the flat-like modifications of the distribution amplitude [21–23], the light-cone holography approach [24, 25], and various model approaches, like a chiral quark model [26] (see also [27–30]), a vector meson dominance model and its modifications [31, 32]. Some other approaches can be found in [33–42].

In this paper we extend and develop the anomaly sum rule approach [7, 8] to the transition form factors with a systematic account of the effects of mixing of η, η' mesons and quark-hadron duality. Also, we performed a new analysis of the anomaly sum rule for π^0 using available experimental data, including the most recent ones from the BELLE collaboration [14].

The paper is organized as follows. In the Section II we give an overview of the anomaly sum rule approach and apply it to analyze the pion (isovector channel of the ASR) and η, η' (octet channel of the ASR) transition form factors. We paid a special attention to the seemingly controversial data from the BABAR [12] and BELLE [14] collaborations. The analysis of different sets of data showed that inclusion of the BABAR data requires a non-(local) OPE correction to the spectral density, while the BELLE data alone neither requires nor excludes it. In the Section III we develop and reformulate the description of mixing, which plays a special role for the η, η' mesons, in a way which does not require the introduction of intermediate non-physical states. The problem of compatibility of different mixing schemes is also discussed. In the Section IV we performed a numerical analysis of the mixing parameters of the $\eta - \eta'$ system basing on the obtained sum rule for the transition form factors. The possibility of the non-OPE correction to the spectral density in the octet channel is investigated as well. The summary is presented in the Section V.

II. ANOMALY SUM RULE APPROACH

The triangle graph amplitude, composed of the axial current $J_{\alpha 5}$ with momentum $p = k + q$ and two vector currents with momenta k and q (VVA amplitude)

$$T_{\alpha\mu\nu}(k, q) = \int d^4x d^4y e^{(ikx+iqy)} \langle 0|T\{J_{\alpha 5}(0) J_\mu(x) J_\nu(y)\}|0\rangle \tag{1}$$

can be decomposed [43] (see also [44, 45]) as

---

[*] Electronic address: klopot@theor.jinr.ru; on leave from Bogolyubov Institute for Theoretical Physics, Kiev, Ukraine
[†] Electronic address: armen@itep.ru
[‡] Electronic address: teryaev@theor.jinr.ru

$$\begin{aligned} T_{\alpha\mu\nu}(k,q) = &F_1\ \varepsilon_{\alpha\mu\nu\rho}k^\rho + F_2\ \varepsilon_{\alpha\mu\nu\rho}q^\rho \\ +\ & F_3\ k_\nu\varepsilon_{\alpha\mu\rho\sigma}k^\rho q^\sigma + F_4\ q_\nu\varepsilon_{\alpha\mu\rho\sigma}k^\rho q^\sigma \\ +\ & F_5\ k_\mu\varepsilon_{\alpha\nu\rho\sigma}k^\rho q^\sigma + F_6\ q_\mu\varepsilon_{\alpha\nu\rho\sigma}k^\rho q^\sigma, \quad (2) \end{aligned}$$

where the coefficients $F_j = F_j(p^2,k^2,q^2;m^2)$, $j = 1,\ldots,6$ are the corresponding Lorentz invariant amplitudes constrained by current conservation and Bose symmetry. Note, that the latter includes the interchange $\mu \leftrightarrow \nu, k \leftrightarrow q$ in the tensor structures and $k^2 \leftrightarrow q^2$ in the arguments of the scalar functions F_j.

In what follows, we consider the case when one of the photons is real ($k^2 = 0$) while the other is real or virtual ($Q^2 = -q^2 \geq 0$). For the invariant amplitude $F_3 - F_6$ the anomaly sum rule (ASR) is given by [5]

$$\int_{4m^2}^{\infty} A_3^{(a)}(s,Q^2;m^2)ds = \frac{1}{2\pi}N_c C^{(a)}\ , \qquad (3)$$

where $A_3 = \frac{1}{2}Im_{p^2}(F_3 - F_6)$, $N_c = 3$ is a number of colors, m is a quark mass and $C^{(a)}$ are the charge factors of components of the axial currents $J_{\alpha 5}^{(a)}$

$$\begin{aligned} J_{\mu 5}^{(3)} &= \frac{1}{\sqrt{2}}(\bar{u}\gamma_\mu\gamma_5 u - \bar{d}\gamma_\mu\gamma_5 d), \\ J_{\mu 5}^{(8)} &= \frac{1}{\sqrt{6}}(\bar{u}\gamma_\mu\gamma_5 u + \bar{d}\gamma_\mu\gamma_5 d - 2\bar{s}\gamma_\mu\gamma_5 s), \\ J_{\mu 5}^{(0)} &= \frac{1}{\sqrt{3}}(\bar{u}\gamma_\mu\gamma_5 u + \bar{d}\gamma_\mu\gamma_5 d + \bar{s}\gamma_\mu\gamma_5 s), \qquad (4) \end{aligned}$$

$$\begin{aligned} C^{(3)} &= \frac{1}{\sqrt{2}}(e_u^2 - e_d^2) = \frac{1}{3\sqrt{2}}, \\ C^{(8)} &= \frac{1}{\sqrt{6}}(e_u^2 + e_d^2 - 2e_s^2) = \frac{1}{3\sqrt{6}}, \\ C^{(0)} &= \frac{1}{\sqrt{3}}(e_u^2 + e_d^2 + e_s^2) = \frac{2}{3\sqrt{3}}. \qquad (5) \end{aligned}$$

Our aim is to investigate the transition form factors of π^0, η and η' mesons, so hereinafter we restrict ourselves to isovector ($a = 3$) and octet ($a = 8$) channels of axial current.

For these channels the ASR (3) has a remarkable property – both perturbative and nonperturbative corrections to the integral are absent [71]. The perturbative corrections are excluded because of Adler-Bardeen theorem [46], while the nonperturbative corrections are also absent as it is expected from 't Hooft's principle. The 't Hooft's principle in its original form [47] implies that the anomalies of the fundamental fields are reproduced on the hadron level. In the dispersive approach this means the absence of the corrections to the dispersive sum rules.

Let us stress, that the spectral density $A_3^{(a)}(s,Q^2;m^2)$ can have both perturbative and nonperturbative corrections (however, the first-order correction $\propto \alpha_s$ is zero in the massless limit [48]), while the integral $\int_{4m^2}^{\infty} A_3^{(a)}(s,Q^2;m^2)ds$ equals exactly $\frac{1}{2\pi}N_c C^{(a)}$.

Saturating the l.h.s. of the three-point correlation function (1) with the resonances and singling out their contributions to the ASR (3) we get the (infinite [7]) sum of resonances with appropriate quantum numbers

$$\pi\sum f_M^a F_{M\gamma} = \int_{4m^2}^{\infty} A_3^{(a)}(s,Q^2;m^2)ds = \frac{1}{2\pi}N_c C^{(a)}. \qquad (6)$$

Here the projections of the axial currents $J_{5\alpha}^{(a)}$ onto one-meson states $M(=\pi^0,\eta,\eta')$ define the coupling (decay) constants f_M^a

$$\langle 0|J_{\alpha 5}^{(a)}(0)|M(p)\rangle = ip_\alpha f_M^a\ , \qquad (7)$$

while the form factors $F_{M\gamma}$ of the transitions $\gamma\gamma^* \to M$ are defined by the matrix elements

$$\int d^4x e^{ikx}\langle M(p)|T\{J_\mu(x)J_\nu(0)\}|0\rangle = \epsilon_{\mu\nu\rho\sigma}k^\rho q^\sigma F_{M\gamma}\ . \qquad (8)$$

The relation (6) expresses the global duality between hadrons and quarks.

A. Isovector channel (π^0)

For a case of the *isovector channel*, the first contribution is given by π^0, while the higher contributions can be absorbed by the "continuum" contribution $\int_{s_0^{(3)}}^{\infty} A_3^{(3)}(s,Q^2;m^2)$, so the ASR (6) takes the form:

$$\pi f_\pi F_{\pi\gamma}(Q^2;m^2) = \frac{1}{2\pi}N_c C^{(3)} - \int_{s_0^{(3)}}^{\infty} A_3^{(3)}(s,Q^2;m^2)ds. \qquad (9)$$

The lower limit $s_0^{(3)}$ of the integral we will refer to as a "continuum threshold", bearing in mind that in a local quark-hadron duality hypothesis it has a meaning of the interval of duality of pion. Also, it can be determined directly from the ASR as it will be demonstrated later.

The spectral density $A_3^{(3)}(s,Q^2;m^2)$ can be calculated from the VVA triangle diagram [5],

$$A_3^{(3)}(s,Q^2;m^2) = \frac{N_c C^{(3)}}{2\pi}\frac{1}{(Q^2+s)^2}\left(Q^2 R + 2m^2\ln\frac{1+R}{1-R}\right), \qquad (10)$$

where $R(s,m^2) = \sqrt{1 - \frac{4m^2}{s}}$, m is a quark mass.

From (9), (10) a straightforward calculation gives an expression for the pion transition form factor,

$$\begin{aligned} F_{\pi\gamma}(Q^2;m^2) &= \frac{1}{2\sqrt{2}\pi^2 f_\pi}\frac{s_0^{(3)}}{s_0^{(3)} + Q^2}\times \\ &\Big[1 - \frac{2m^2}{s_0^{(3)}}\big(\frac{2}{R_0+1} + \ln\frac{1+R_0}{1-R_0}\big)\Big], \qquad (11) \end{aligned}$$

where $R_0 = R(s_0^{(3)}, m^2)$. This expression (to our best knowledge, for the first time) takes into account the contribution of quark mass. However, in what follows, we neglect the contributions of all light (u, d, s) quark masses to the ASR in both isovector and octet channels. The phenomenological applications of the quark mass effects will be considered in a separate paper.

In this approximation, the spectral density (10) is proportional to $\delta(s)$ at $Q^2 = 0$, so the continuum term in the ASR (9) goes to zero. This corresponds to the fact that contributions of axial states are zero at $Q^2 = 0$, and contributions of higher pseudoscalar states should be suppressed in order for the axial current to conserve in the chiral limit.

Relying on the local quark-hadron duality hypothesis, the analysis of the two-point correlation function gives the value for the continuum threshold $s_0^{(3)} = 0.75$ GeV^2 [49]. Actually, $s_0^{(3)}$ can be determined directly from the high-Q^2 asymptotic of ASR, where the QCD factorization predicts the value of transition form factor $Q^2 F^{as}_{\pi\gamma} = \sqrt{2} f_\pi$ [15]. The high-Q^2 limit of (11) $(m = 0)$ immediately leads to $s_0^{(3)} = 4\pi^2 f_\pi^2 = 0.67$ GeV^2. This expression, substituted to (11) with $m = 0$ gives

$$F_{\pi\gamma}(Q^2; 0) = \frac{1}{2\sqrt{2}\pi^2 f_\pi} \frac{4\pi^2 f_\pi^2}{4\pi^2 f_\pi^2 + Q^2}, \tag{12}$$

so it proves the Brodsky-Lepage interpolation formula for the pion transition form factor [50], which was later confirmed by Radyushkin [51] in the approach of local quark-hadron duality. Let us stress, that in this way we found that it is a direct consequence of the anomaly sum rule (which is an exact nonperturbative QCD relation).

Currently the data from the CELLO [52], CLEO [53], BABAR [12] and BELLE [14] collaborations cover the region of $Q^2 = 0.7 - 35$ GeV^2 (see Fig. 1). While at $Q^2 < 10$ GeV^2 they are consistent, at larger virtualities the BABAR and a newly released BELLE data are quite different. In this situation we will consider two sets of data: CELLO+CLEO+BELLE (I) and CELLO+CLEO+BABAR (II).

When compared to the experimental data set (I), Eq. (12) gives a reasonable, consistent with the data description ($\chi^2/d.o.f. = 1.01, d.o.f = 35$, see the dashed line in Fig. 1). For the data set (II) the description is worse ($\chi^2/d.o.f. = 2.29, d.o.f. = 37$). So, if the data of the BABAR collaboration are correct, we come to a violation of the ASR-based expression for $F_{\pi\gamma}$ (12). This means that the spectral density (10) must have a substantial correction δA_3, which results in corrections to the continuum $\delta I_{cont} = \int_{s_0^{(3)}}^{\infty} \delta A_3 ds$ and pion $\delta I_\pi = \int_{4m^2}^{s_0^{(3)}} \delta A_3 ds$ contributions. At the same time, as the *full* integral remains constant, the corrections can be related:

$$\delta I_\pi + \delta I_{cont} = 0. \tag{13}$$

It is important, that the main terms of the continuum I_{cont} and the pion I_π contributions have essentially different Q^2 behaviour

$$I_{cont} = \int_{s_0^{(3)}}^{\infty} A_3^{(3)}(s, Q^2) ds = \frac{1}{2\sqrt{2}\pi} \frac{Q^2}{s_0^{(3)} + Q^2}, \tag{14}$$

$$I_\pi = \int_0^{s_0^{(3)}} A_3^{(3)}(s, Q^2) ds = \frac{1}{2\sqrt{2}\pi} \frac{s_0^{(3)}}{s_0^{(3)} + Q^2}, \tag{15}$$

so the $1/Q^2$ power correction to the continuum contribution is of the order of the main term of the pion contribution.

Let us now discuss the sources of possible corrections to the spectral densities which in our approach are the counterparts of the NLO and higher twist corrections in the accurate pQCD fits [18]. Note that one-loop corrections to the spectral densities of all structures in the VVA correlator in the massless case are zero, which can be easily deduced from the results of [48]. If this nullification is due to conformal invariance [54, 55], one may expect the two- and higher loop corrections to be nonzero due to the beta-function effects, which is recently observed in the soft-photon approximation [56] . Nevertheless, these higher α_s corrections as well as the local OPE-induced corrections are small enough to produce an enhancement in pion transition form factor shown by BABAR: $\delta F_{\pi\gamma} \sim log(Q^2)/Q^2$. Clearly, from dimensional arguments, such term cannot appear from the local OPE [7] . Thus, to comply with the ASR, the correction should be of a non-(local) OPE origin simulating the contribution of the operator of dimension 2. Among the possible sources of such correction are non-local condensates, instantons or short strings [57].

Although the exact form of such correction is not yet known, we can construct the simplest form of it relying on general requirements. Namely, the correction should vanish at $s_0^{(3)} \to \infty$ (the continuum contribution vanishes), at $s_0^{(3)} \to 0$ (the full integral has no corrections), at $Q^2 \to \infty$ (the perturbative theory works at large Q^2) and at $Q^2 \to 0$ (anomaly perfectly describes pion decay width). Supposing the correction contains rational functions and logarithms of Q^2, the simplest form of the correction satisfying those limits is

$$\delta I_\pi = \frac{1}{2\sqrt{2}\pi} \frac{\lambda s_0^{(3)} Q^2}{(s_0^{(3)} + Q^2)^2} (\ln \frac{Q^2}{s_0^{(3)}} + \sigma), \tag{16}$$

where λ and σ are dimensionless parameters. Then the pion transition form factor with this correction reads:

$$F_{\pi\gamma}(Q^2) = \frac{1}{\pi f_\pi}(I_\pi + \delta I_\pi) =$$
$$\frac{1}{2\sqrt{2}\pi^2 f_\pi} \frac{s_0^{(3)}}{s_0^{(3)} + Q^2} \Big[1 + \frac{\lambda Q^2}{s_0^{(3)} + Q^2} (\ln \frac{Q^2}{s_0^{(3)}} + \sigma)\Big]. \tag{17}$$

For the continuum threshold we will use $s_0^{(3)} = 4\pi^2 f_\pi^2 = 0.67$ GeV^2, which implies that at very high Q^2 the factorization restores. Nevertheless, if the factorization is violated at all Q^2, one can use a different value for $s_0^{(3)}$. Moreover, one can even consider the dependence of $s_0^{(3)}$ on Q^2 [38], which can lead to an effective change of σ.

The fit of (17) to the data set (II) gives $\lambda = 0.14$, $\sigma = -2.36$ with $\chi^2/d.o.f. = 0.94$ $(d.o.f. = 35)$. The plot of $Q^2 F_{\pi\gamma}$ for these parameters is depicted in Fig. 1 as a solid curve. Note, that the proposed correction changes its sign at $Q^2 \approx 8$ GeV^2 improving the description in both regions: at $Q^2 \lesssim 8$ GeV^2 and at $Q^2 \gtrsim 8$ GeV^2. Also, $F_{\pi\gamma}$ (17) with these parameters λ, σ (obtained from the fit to the data (II)) describes well also the data set (I) with $\chi^2/d.o.f. = 0.84$ $(d.o.f. = 35)$.

The summary of the fitting results for $F_{\pi\gamma}$ with (Eq. (17)) and without (Eq. (12)) correction for different sets of data are shown in Table I. One can see, that the data sets involving the BABAR data require taking into account the correction. The data sets which do not involve the BABAR data may be described without such correction, although the correction may improve description of the data.

Let us emphasize, that the correction (16) requires a $\log Q^2$ term in δA_3 itself. This form of the correction is different from the one proposed in [10] and could match it only if the pre-logarithmic factor in (16) did not depend on $s_0^{(3)}$. However, such factor would violate the mentioned above requirement of nullification of the correction (in the limit $s_0^{(3)} \to 0$).

Also, there is a clear distinction with the natural emergence of the $\log Q^2$ term in the triangle amplitude (which was used for description of the BABAR data), where triangle amplitude itself is used as a model for pion transition form factor [26]. Such a description is based on the consideration of the triangle diagram as a model for the transition form factors and applies the PCAC relation for the matrix elements involving large virtualities. To our opinion this procedure is not justified to the same degree of rigor as for the soft processes.

It is interesting to note, that extending the expression for the pion transition form factor (11) into the timelike region, one immediately gets a pole at $Q^2 = s_0^{(3)}$ which is numerically close to the mass of ρ meson. This can be a kind of interplay between anomaly and vector dominance.

B. Octet channel (η, η')

For a case of the *octet channel*, where the main contributions are from η and η' mesons, treating the ASR in the same way as for the isovector channel leads to

$$f_\eta^8 F_{\eta\gamma}(Q^2) + f_{\eta'}^8 F_{\eta'\gamma}(Q^2) = \frac{1}{2\sqrt{6}\pi^2} \frac{s_0^{(8)}}{s_0^{(8)} + Q^2}. \qquad (18)$$



FIG. 1: (color online). Pion transition form factor (multiplied by Q^2) with correction (Eq. (17), solid curve) and without correction (Eq. (12), dashed curve) as a function of Q^2 compared with experimental data.

In comparison with the isovector channel, here we have some differences. Due to significant mixing in $\eta - \eta'$ system the η' meson has a significant contribution to the octet channel. Since η' decays into two real photons, it should be taken into account explicitly along with the η meson.

Let us note, that in the octet channel, the reliable estimation of $s_0^{(8)}$ from the usual QCD sum rule method meets difficulties (see e.g. discussion in [9]). Fortunately, the ASR approach allows to determine $s_0^{(8)}$ in the octet channel from high-Q^2 asymptotic, just the same way as in the isovector channel. Generalization of the pion case gives the asymptotes for the η, η' transition form factors [58, 59] $(M = \eta, \eta')$:

$$Q^2 F_{M\gamma}^{as} = 6(C^{(8)} f_M^8 + C^{(0)} f_M^0). \qquad (19)$$

So, the $Q^2 \to \infty$ limit of the ASR (18) gives

$$s_0^{(8)} = 4\pi^2((f_\eta^8)^2 + (f_{\eta'}^8)^2 + 2\sqrt{2}[f_\eta^8 f_\eta^0 + f_{\eta'}^8 f_{\eta'}^0]). \qquad (20)$$

In the $Q^2 = 0$ limit the transition form factors are expressed in terms of the two-photon decay widths of mesons, so the ASR (18) takes the form:

$$f_\eta^8 F_{\eta\gamma}(0) + f_{\eta'}^8 F_{\eta'\gamma}(0) = \frac{1}{2\sqrt{6}\pi^2} \frac{s_0^{(8)}}{s_0^{(8)} + Q^2}, \qquad (21)$$

where

$$F_{M\gamma}(0) = \sqrt{\frac{4\Gamma_{M\to\gamma\gamma}}{\pi\alpha^2 m_M^3}}.$$

Solving equations (20), (21) with respect to $s_0^{(8)}$ and one of the decay constants f_M^a and substituting them into general ASR (18) we can relate the transition form factors $F_{\eta'\gamma}(Q^2)$ and the decay constants f_M^a.

	$\delta I_\pi = 0:$ $\chi^2/d.o.f.$	$\delta I_\pi \neq 0:$ $\chi^2/d.o.f.$	λ	σ
CELLO+CLEO+BABAR+BELLE	1.86	0.91	0.12	-2.50
CELLO+CLEO+BELLE (set I)	1.01	0.46	0.07	-3.03
CELLO+CLEO+BABAR (set II)	2.29	0.94	0.14	-2.36
BABAR	3.61	0.99	0.20	-2.39
BELLE	0.80	0.40	0.14	-2.86

TABLE I: $\chi^2/d.o.f.$ obtained for Eq. (12) ($\delta I_\pi = 0$) and the best fits of Eq. (17) ($\delta I_\pi \neq 0$) to different data sets.

Actually, the four decay constants f_M^a can be related basing on a particular mixing scheme. In [8] the ASR was analyzed for several sets of parameters and mixing schemes. The more general consideration of mixing schemes and extraction of the mixing parameters are performed in the next sections.

III. MIXING

The problem of mixing in the $\eta-\eta'$ system is usually addressed either in the octet-singlet ($SU(3)$) or quark-flavor mixing scheme (see e.g. [60, 61] and references therein). Basically, meson mixing implies that the "non-diagonal" decay constants $f_M, M=\eta,\eta'$ in Eq. (7) are non-zero. The π^0 and the isovector current can be decoupled from the $\eta-\eta'$ system and octet and singlet currents because of a very small mixing.

Let us recall the common approach to the mixing, when physical states are represented as a linear combination of states with definite $SU(3)_f$ quantum numbers or quark-flavor content. But, since these states do not have definite masses, one cannot write the analogue of Eq. (7) with these states instead of physical states η, η'. Indeed, if a state has a definite momentum p^μ it also has a definite mass $m^2 = p_\mu p^\mu$.

One can avoid this problem if formulates the mixing in terms of the fields ϕ_i related to the physical states $|i\rangle$ by the matrix elements $\langle 0|\phi_i|j\rangle = \delta_{ij}$.

It is well known, that the field of the pion ϕ_π is defined from the divergence of the isovector component of axial channel as a PCAC relation,

$$\partial_\mu J^{(3)}_{\mu 5} = f^{(3)}_\pi m^2_\pi \phi_\pi. \qquad (22)$$

To consider the η and η' mixing, one can write down a straightforward generalization of this relation,

$$\partial_\mu \mathbf{J}_{\mu 5} = \mathbf{F}\mathbf{M}\mathbf{\Phi}, \qquad (23)$$

where we introduced the matrix notations:

$$\mathbf{J}_{\mu 5} \equiv \begin{pmatrix} J^\alpha_{\mu 5} \\ J^\beta_{\mu 5} \end{pmatrix}, \mathbf{F} \equiv \begin{pmatrix} f^\alpha_\eta & f^\alpha_{\eta'} & f^\alpha_G & \cdots \\ f^\beta_\eta & f^\beta_{\eta'} & f^\beta_G & \cdots \end{pmatrix}, \mathbf{\Phi} \equiv \begin{pmatrix} \phi_\eta \\ \phi_{\eta'} \\ \phi_G \\ \vdots \end{pmatrix},$$
$$\mathbf{M} \equiv diag(m^2_\eta, m^2_{\eta'}, m^2_G, \cdots). \qquad (24)$$

The vector $\mathbf{J}_{\mu 5}$ in the l.h.s. consists of the components of the axial current of definite $SU(3)$ symmetry, a so-called octet-singlet basis ($\alpha = 8, \beta = 0$, see Eq. (4)), or of the components of the axial current with the decoupled light and strange quark composition, a so-called quark-flavor basis ($\alpha = q, \beta = s$):

$$J^q_{\mu 5} = \frac{1}{\sqrt{2}}(\bar{u}\gamma_\mu\gamma_5 u + \bar{d}\gamma_\mu\gamma_5 d), \; J^s_{\mu 5} = \bar{s}\gamma_\mu\gamma_5 s. \qquad (25)$$

The elements of matrix $\mathbf{F}$ are the meson decay constants defined in (7). The vector $\mathbf{\Phi}$ of physical fields contains the fields of η and η' mesons ϕ_η and $\phi_{\eta'}$ and the fields of higher mass states, which we denote as $\phi_G, \ldots$.

It is possible and commonly to get an additional model constraint for the matrix $\mathbf{F}$ which is fulfilled by applying the respective mixing scheme. Let us first introduce the new vector of fields $\widetilde{\mathbf{\Phi}}$ relating the $SU(3)$ symmetry property or the quark-flavor contents of currents and meson fields. The first two components of $\widetilde{\mathbf{\Phi}}$ are labelled with the same indices α and β as the currents and correspond to the same symmetry ($\alpha = 8, \beta = 0$) or quark content ($\alpha = q, \beta = s$). The relation between $\mathbf{\Phi}$ and $\widetilde{\mathbf{\Phi}}$ is provided by the orthogonal transformation $\mathbf{U}$

$$\widetilde{\mathbf{\Phi}} = \mathbf{U}\mathbf{\Phi}, \;\; \widetilde{\mathbf{\Phi}} \equiv \begin{pmatrix} \tilde{\phi}_\alpha \\ \tilde{\phi}_\beta \\ \tilde{\phi}_G \\ \vdots \end{pmatrix}. \qquad (26)$$

In terms of these fields the Eq. (23) can be rewritten as

$$\partial_\mu \mathbf{J}_{\mu 5} = \widetilde{\mathbf{F}}\widetilde{\mathbf{M}}\widetilde{\mathbf{\Phi}}, \qquad (27)$$

where $\widetilde{\mathbf{F}} = \mathbf{F}\mathbf{U}$, $\widetilde{\mathbf{M}} = \mathbf{U}^\mathbf{T}\mathbf{M}\mathbf{U}$.

In our notations the octet-singlet (quark-flavor) mixing scheme implies that the matrix $\widetilde{\mathbf{F}}$ has a (rectangular) diagonal form in the respective octet-singlet (quark-flavor) basis,

$$\widetilde{\mathbf{F}} = \begin{pmatrix} f_\alpha & 0 & 0 & \ldots \\ 0 & f_\beta & 0 & \ldots \end{pmatrix}. \qquad (28)$$

This relation can be obtained from the effective lagrangian $\mathcal{L}$ which contains an interaction term $\Delta\mathcal{L}_{int} = \frac{1}{2}\widetilde{\mathbf{\Phi}}^\mathbf{T}\widetilde{\mathbf{M}}\widetilde{\mathbf{\Phi}} = \frac{1}{2}\sum\limits_{i,j}\tilde{m}^2_{ij}\tilde{\phi}_i\tilde{\phi}_j$:

$$\partial_\mu J^a_{\mu 5} = f_a \frac{\delta\mathcal{L}}{\delta\tilde{\phi}_a} = f_a \sum_k \tilde{m}^2_{ak}\tilde{\phi}_k, \; a = \alpha, \beta. \qquad (29)$$

Note, that from the requirement that matrix $\mathbf{FU}$ has a (rectangular) diagonal form (28) immediately follows that $\mathbf{FF^T}$ is a diagonal matrix. So, imposing the mixing scheme is equivalent to imposing the constraint for the decay constants:

$$f^{\alpha}_{\eta} f^{\beta}_{\eta} + f^{\alpha}_{\eta'} f^{\beta}_{\eta'} + f^{\alpha}_{G} f^{\beta}_{G} + ... = 0. \tag{30}$$

Here the sum is over all physical meson states included in the vector $\mathbf{\Phi}$.

If we restrict ourselves to consideration of the η and η' mesons only, then the decay constants form a 2x2 matrix and in the octet-singlet and quark-flavor bases satisfy the respective diagonality constraints,

$$f^{8}_{\eta} f^{0}_{\eta} + f^{8}_{\eta'} f^{0}_{\eta'} = 0, \tag{31}$$

$$f^{q}_{\eta} f^{s}_{\eta} + f^{q}_{\eta'} f^{s}_{\eta'} = 0. \tag{32}$$

For instance, in the case of the octet-singlet mixing scheme (31) the matrix of decay constants can be expressed in terms of one mixing angle θ and two parameters f_8, f_0, forming the well-known one-angle mixing scheme:

$$\mathbf{F_{80}} = \begin{pmatrix} f^{8}_{\eta} & f^{8}_{\eta'} \\ f^{0}_{\eta} & f^{0}_{\eta'} \end{pmatrix} = \begin{pmatrix} f_8 \cos\theta & f_8 \sin\theta \\ -f_0 \sin\theta & f_0 \cos\theta \end{pmatrix}. \tag{33}$$

Similarly, if the quark-flavor mixing scheme restriction (32) is applied, then it is common to express the decay constants in terms of parameters ϕ, f_8, f_0,

$$\mathbf{F_{qs}} = \begin{pmatrix} f^{q}_{\eta} & f^{q}_{\eta'} \\ f^{s}_{\eta} & f^{s}_{\eta'} \end{pmatrix} = \begin{pmatrix} f_q \cos\phi & f_q \sin\phi \\ -f_s \sin\phi & f_s \cos\phi \end{pmatrix}. \tag{34}$$

While either of the mixing schemes (octet-singlet or quark-flavor) is self-consistent, they are incompatible [9, 60]. Indeed, octet-singlet and quark-flavor bases of axial currents are related by means of rotation matrix:

$$\begin{pmatrix} J^{8}_{\mu 5} \\ J^{0}_{\mu 5} \end{pmatrix} = \mathbf{V}(\alpha) \begin{pmatrix} J^{q}_{\mu 5} \\ J^{s}_{\mu 5} \end{pmatrix}, \; \mathbf{V}(\alpha) = \begin{pmatrix} \cos\alpha & -\sin\alpha \\ \sin\alpha & \cos\alpha \end{pmatrix}, \tag{35}$$

where $\tan\alpha = \sqrt{2}$. Then, as follows from (23), the matrices of decay constants $\mathbf{F}_{\alpha\beta} \equiv \mathbf{F}$ (24) in the octet-singlet (33) and quark-flavor (34) bases are related as

$$\mathbf{F_{80}} = \mathbf{V}(\alpha) \mathbf{F_{qs}}, \tag{36}$$

and so

$$\mathbf{F_{80}} \mathbf{F^T_{80}} = \mathbf{V}(\alpha) \mathbf{F_{qs}} \mathbf{F^T_{qs}} \mathbf{V}(\alpha)^{\mathbf{T}}. \tag{37}$$

We see that in general case the decay constants cannot follow the octet-singlet and quark-flavor mixing scheme simultaneously (since the matrices $\mathbf{F_{80}F^T_{80}}$ and $\mathbf{F_{qs}F^T_{qs}}$ cannot be diagonal simultaneously). The bases are compatible only if $f_8 = f_0$ ($f_q = f_s$), i.e. in case of the exact $SU(3)_f$ symmetry.

Although the octet-singlet mixing scheme is more convenient for the anomaly sum rule relations, which are exact in case of isovector and octet channels, there are arguments from chiral perturbation theory against it [62, 63] (see also [60, 64]).

The mixing scheme-independent extraction of the decay constants from the experimental data can finally tell us which basis is more adequate for describing the mixing in the $\eta - \eta'$ system. This problem will be addressed in the next section.

IV. OCTET CHANNEL: NUMERICAL ANALYSIS

In the Section II, as a consequence of the ASR in the octet channel, we obtained the relation between transition form factors and decay constants of η and η' mesons. In this section we use this relation to analyze the decay constants in different mixing schemes, described in the Sec. III.

First, let us consider the *octet-singlet mixing scheme.* In order to determine the mixing parameters of this scheme, we employ the mixing scheme constraint (31) and the ASR relations (18), (20), (21). In terms of the parameters (33) the Eq. (20) reads: $s^{(8)}_0 = 4\pi^2 f^2_8$. Then, the regions in f_8, θ parameter space which are constrained by the fit of Eq. (18) ($\chi^2/d.o.f. < 1$) to the BABAR data [13] and by the Eq. (21) (the experimental errors of $\Gamma_{\eta(\eta')\to 2\gamma}$ are taken into account) are shown in Fig. 2. The yellow intersection determines the parameters, which can be estimated as: $f_8 = (0.88 \pm 0.04) f_\pi, \theta = -(14.2 \pm 0.7)^\circ$.

In order to determine the constant f_0 (which does not enter the Eqs. (18), (20),(21) in case of the octet-singlet mixing scheme) we need an additional constraint.

As an additional constraint it is convenient to use the ratio of radiative decays of J/Ψ for which the more cumbersome contribution of the gluonic anomaly is under control. Indeed, according to [66], the radiative decays $J/\Psi \to \eta(\eta')\gamma$ are dominated by the non-perturbative gluonic matrix elements, and the ratio of the decay rates $R_{J/\Psi} = (\Gamma(J/\Psi) \to \eta'\gamma)/(\Gamma(J/\Psi) \to \eta\gamma)$ is given by

$$R_{J/\Psi} = \left| \frac{\langle 0 \mid G\widetilde{G} \mid \eta' \rangle}{\langle 0 \mid G\widetilde{G} \mid \eta \rangle} \right|^2 \left(\frac{p_{\eta'}}{p_\eta} \right)^3, \tag{38}$$

where $p_{\eta(\eta')} = M_{J/\Psi}(1 - m^2_{\eta(\eta')}/M^2_{J/\Psi})/2$.

The divergencies of the octet and singlet components of the axial current in terms of quark fields are



FIG. 2: (color online). Octet-singlet mixing scheme parameters f_8, θ. Blue region – constraint of Eq. (18) ($\chi^2/d.o.f. < 1$), red region – constraint of Eq. (21) (experimental uncertainties are taken into account).

$$\partial_\mu J^8_{\mu 5} = \frac{1}{\sqrt{6}}(m_u \overline{u}\gamma_5 u + m_d \overline{d}\gamma_5 d - 2m_s \overline{s}\gamma_5 s), \tag{39}$$

$$\partial_\mu J^0_{\mu 5} = \frac{1}{\sqrt{3}}(m_u \overline{u}\gamma_5 u + m_d \overline{d}\gamma_5 d + m_s \overline{s}\gamma_5 s) + \frac{1}{2\sqrt{3}}\frac{3\alpha_s}{4\pi}G\widetilde{G}. \tag{40}$$

Taking matrix elements of the above expressions between vacuum and $\eta(\eta')$ states and neglecting the u and d quark masses, the ratio (38) can be expressed [67] in terms of the decay constants (7) as follows:

$$R_{J/\Psi} = \left(\frac{f^8_{\eta'} + \sqrt{2}f^0_{\eta'}}{f^8_{\eta} + \sqrt{2}f^0_{\eta}}\right)^2 \left(\frac{m_{\eta'}}{m_\eta}\right)^4 \left(\frac{p_{\eta'}}{p_\eta}\right)^3. \tag{41}$$

The current experimental value of this ratio is $R_{J/\Psi} = 4.67 \pm 0.15$ [65].

Employing this ratio for the octet-singlet mixing scheme and taking into account Eqs. (18) and (21) one can determine the singlet constant: $f_0 = (0.81 \pm 0.07) f_\pi$. So, the full set of constants of the octet-singlet scheme is:

$$\begin{aligned} f_8 &= (0.88 \pm 0.04) f_\pi, \\ f_0 &= (0.81 \pm 0.07) f_\pi, \\ \theta &= -(14.2 \pm 0.7)^\circ. \end{aligned} \tag{42}$$

For the *quark-flavour mixing scheme* we can perform the similar analysis, using the constraint of the scheme (32), Eqs. (20), (21), (41) and fitting the ASR (18) to the BABAR data [13]. The decay constants of quark-flavour basis $f^{q,s}_{\eta,\eta'}$ are expressed in terms of those of octet-singlet basis $f^{8,0}_{\eta,\eta'}$ by means of the Eq. (36). In terms of the mixing parameters f_q, f_s, ϕ (34) the Eqs. (20), (41) read:



FIG. 3: (color online). Quark-flavour mixing scheme parameters f_q, f_s. Blue region – constraint of Eq. (18) ($\chi^2/d.o.f. < 1$), red region – constraint of Eq. (21) (experimental uncertainties are taken into account).

$$s_0^{(8)} = (4/3)\pi^2(5f_q^2 - 2f_s^2), \tag{43}$$

$$R_{J/\Psi} = (\tan\phi)^2 \left(\frac{m_{\eta'}}{m_{\eta'}}\right)^4 \left(\frac{p_{\eta'}}{p_\eta}\right)^3. \tag{44}$$

The Eq. (44) determines the parameter $\phi = (38.1 \pm 0.5)^\circ$. Then the other two parameters f_s, f_q can be estimated from the Eqs. (21) and (18). The plot of the regions constrained by these equations is shown in Fig. 3. The red band indicates the constraint of the Eq. (21) (experimental errors of $R_{J/\Psi}, \Gamma_{\eta(\eta')\to 2\gamma}$ are taken into account) and the blue band indicates the fit of the Eq. (18) to the BABAR data at $\chi^2/d.o.f. < 1$ level. One can observe two regions where both equations are compatible. We have chosen the physically motivated one, where $f_q, f_s > f_\pi$. So, the yellow intersection in Fig. 3 determines the parameters for the quark-flavour mixing scheme, which give the following ranges for them:

$$\begin{aligned} f_q &= (1.20 \pm 0.15) f_\pi, \\ f_s &= (1.65 \pm 0.25) f_\pi, \\ \phi &= (38.1 \pm 0.5)^\circ. \end{aligned} \tag{45}$$

The obtained mixing parameters (45) are in agreement with those obtained in other approaches [60, 68, 69].

It is interesting also to get the *mixing scheme independent* constraints on the decay constants. Equations (21), (41) allow to exclude two of the four decay constants which enter the ASR (18). The continuum threshold $s_0^{(8)}$ is excluded using (20). In Fig. 4 the levels of $\chi^2/d.o.f.$ function of Eq. (18) in the space of



FIG. 4: (color online). Independent (of mixing scheme) estimation of decay constants. Black thin curve is a $\chi^2/d.o.f. = 1$ level, red dot is a minimum of χ^2 of Eq. (18). Green solid and orange dashed lines indicate the constraints of the octet-singlet (31) and quark-flavor (32) mixing schemes respectively.

constants f_η^8, f_η^0 are shown (BABAR experimental data on $F_{\gamma\eta(\eta')}(Q^2)$ and mean values of $R_{J/\Psi}, \Gamma_{\eta(\eta')\to 2\gamma}$ are used). One can see, that $\chi^2/d.o.f. < 1$ requirement (black curve) allows a rather wide range of the parameters. However, the minimum of χ^2 is reached at the point $f_\eta^8 = 1.11 f_\pi, f_\eta^0 = 0.16 f_\pi$ ($\chi^2/d.o.f. = 0.84$, indicated as a red dot). Then (21), (41) allow to determine the other two constants: $f_{\eta'}^8 = -0.42 f_\pi, f_{\eta'}^0 = 1.04 f_\pi$. Therefore, the full set of decay constants of the mixing scheme-independent extraction is:

$$\begin{pmatrix} f_\eta^8 & f_{\eta'}^8 \\ f_\eta^0 & f_{\eta'}^0 \end{pmatrix} = \begin{pmatrix} 1.11 & -0.42 \\ 0.16 & 1.04 \end{pmatrix} f_\pi. \qquad (46)$$

The constraints of the octet-singlet (green solid curve) (31) and quark-flavor (orange dashed curve) (32) are also depicted in Fig. 4. We see that both considered mixing schemes are consistent with the scheme-independent analysis based on the ASR at the level of $\chi^2/d.o.f. < 1$, even if other experimental errors are not taken into account. However, the least χ^2 is reached in the region lying outside of both mixing scheme curves. Further improvement of the experimental data can clear up the question of validity of different schemes and give more precise values of the mixing parameters.

One can expect that the possible non-OPE correction to the spectral density, discussed for the isovector channel, should manifest in the same way in the octet channel also. Although the BABAR data [13] do not show such a sturdy growth of $Q^2F_{\eta\gamma}$ and $Q^2F_{\eta'\gamma}$ as they do for $Q^2F_{\pi\gamma}$, the octet combination of the transition form factors $Q^2(f_\eta^8 F_{\eta\gamma} + f_{\eta'}^8 F_{\eta'\gamma})$ does reveal a possible growth [72].

Expecting the similarity of the correction to the spectral density in the isovector and octet channels, we suppose that the correction in the octet channel has the same form as (16),



FIG. 5: (color online). ASR with correction ((48), solid line) and without correction ((18), dashed line) for the octet-singlet mixing scheme parameters (42) compared with experimental data (blue points – CLEO, red points – BABAR).

$$\delta I_8 = -\int_{s_0^{(8)}}^{\infty} \delta A_3^{(8)} ds = \frac{1}{2\sqrt{6}\pi} \frac{\lambda s_0^{(8)} Q^2}{(s_0^{(8)}+Q^2)^2} (\ln \frac{Q^2}{s_0^{(8)}} + \sigma). \qquad (47)$$

This correction results in an additional term $\delta I_8/\pi$ in the r.h.s. of (18),

$$f_\eta^8 F_{\eta\gamma}(Q^2) + f_{\eta'}^8 F_{\eta'\gamma}(Q^2) = \frac{1}{2\sqrt{6}\pi^2} \frac{s_0^{(8)}}{s_0^{(8)}+Q^2} + \frac{1}{2\sqrt{6}\pi^2} \frac{\lambda s_0^{(8)} Q^2}{(s_0^{(8)}+Q^2)^2} (\ln \frac{Q^2}{s_0^{(8)}} + \sigma). \qquad (48)$$

The plot of the Eq. (48) (solid line) and Eq. (18) (dashed line) for different mixing schemes are shown in Figs. 5,6,7. The parameters $\lambda = 0.14, \sigma = -2.36$ are taken the same as obtained for the pion case (data set II), the decay constants (central values) for different mixing schemes are employed from (42), (45) and (46). We see, that the current precision of the experimental data on the η and η' transition form factors could accommodate the same correction as in the pion case, however does not require it.

Finally, let us make the following remark. In this paper we developed the approach which does not rely on introduction of the non-physical states which have no definite masses. At the same time, in literature is widely discussed a hypothesis (see e.g. [13, 70]) that the transition form factor of the non-physical state $|q\rangle \equiv \frac{1}{\sqrt{2}}(|\bar{u}u\rangle + |\bar{d}d\rangle)$ is related to the pion form factor as $F_{q\gamma}(Q^2) = (5/3)F_{\pi\gamma}(Q^2)$ (where the numerical factor comes from the quark charges: $(e_u^2+e_d^2)/(e_u^2-e_d^2) = 5/3$). The states $|q\rangle$ and $|s\rangle \equiv |\bar{s}s\rangle$ are assumed to be expressed in terms of the physical states $|\eta\rangle$, $|\eta'\rangle$ via the quark-flavor mixing scheme [73],

$$|q\rangle = \cos\phi|\eta\rangle + \sin\phi|\eta'\rangle, |s\rangle = -\sin\phi|\eta\rangle + \sin\phi|\eta'\rangle. \quad (49)$$



FIG. 6: (color online). ASR with correction ((48), solid line) and without correction ((18), dashed line) for the quark-flavor mixing scheme parameters (45) compared with experimental data (blue points – CLEO, red points – BABAR).



FIG. 7: (color online). ASR with correction ((48), solid line) and without correction ((18), dashed line) for the mixing scheme-independent parameters (46) compared with experimental data (blue points – CLEO, red points – BABAR).

Then one can relate the form factors:

$$\frac{5}{3}F_{\pi\gamma} = F_{\eta\gamma}\cos\phi + F_{\eta'\gamma}\sin\phi. \tag{50}$$

Let us now try to incorporate this hypothesis into our approach. For this purpose, combining (18), (50) and using (11), (36) we can express the η, η' transition form factors in terms of the constants f_q, f_s, ϕ,

$$F_{\eta\gamma}(Q^2) = \frac{1}{4\pi^2 f_s f_\pi}\frac{s_0^{(3)}(\sqrt{2}f_s\cos\phi - f_q\sin\phi)}{s_0^{(3)}+Q^2} + \frac{1}{4\pi^2 f_s}\frac{s_0^{(8)}\sin\phi}{s_0^{(8)}+Q^2}, \tag{51}$$

$$F_{\eta'\gamma}(Q^2) = \frac{1}{4\pi^2 f_s f_\pi}\frac{s_0^{(3)}(\sqrt{2}f_s\sin\phi + f_q\cos\phi)}{s_0^{(3)}+Q^2} + \frac{1}{4\pi^2 f_s}\frac{s_0^{(8)}\cos\phi}{s_0^{(8)}+Q^2}, \tag{52}$$

where $s_0^{(3)} = 4\pi^2 f_\pi^2$, $s_0^{(8)} = (4/3)\pi^2(5f_q^2 - 2f_s^2)$.



FIG. 8: (color online). Combinations $F_{\eta\gamma}Q^2$ (blue solid line) and $F_{\eta'\gamma}Q^2$ (red dashed line) from Eqs. (51) and (52) respectively as functions of Q^2 compared with experimental data.

The plot of Eqs. (51), (52) with constants from our analysis (45) $f_q = 1.20 f_\pi, f_s = 1.65 f_\pi, \phi = 38.1^\circ$ in comparison with experimental data is shown in Fig. 8. One can observe a reasonably good agreement with the experimental data. For the decay constants of Ref. [60] one gets a good description also.

The agreement with the experimental data may indicate that the effect of strong anomaly for $\frac{1}{\sqrt{2}}|\bar{u}u + \bar{d}d\rangle$ state is small and the strong anomaly predominantly appears in the $\bar{s}s$ channel. This statement can be rigorously checked by means of the anomaly sum rule for the singlet channel, which we postpone to future work.

V. SUMMARY

The exact anomaly sum rule allows to derive an expression for the pion transition form factor at arbitrary Q^2, giving the proof for the Brodsky-Lepage interpolation formula. At $Q^2 = 0$ it is related to the pion decay width, while at large Q^2, basing on the factorization approach, it allows to define the pion interval of duality which numerically appears to be close to the value defined from the two-point correlator sum rule analysis. However, the proposed approach may be applied even when the factorization is broken. It was exactly the situation supported by BABAR data requiring a small nonperturbative correction [7] to the continuum spectral density. Having dimension 2, it cannot appear in (a local) OPE and should be attributed to, say, instantons or short strings.

In this work we included in our analysis the recent BELLE collaboration data. The main conclusions are the following. Although the BELLE data themselves may be described without the mentioned correction, but they do not also exclude its possibility. Unless the BABAR data will be disproved, the need for the correction remains. This is supported by the Table 1, where the fits for various combinations of the data are shown.

In the generalization of our approach to the η and η' mesons, the data may be described without such a correction. However, the possibility of the correction, similar to the discussed for the pion case is not excluded by the current experimental data and supported by the slight growth of the octet combination of transition form factors. So we can conclude, that the correction to the spectral density, first introduced in [7] seems to be universal for both, isovector and octet channels. This conclusion is in an agreement with a recent discussion in [11].

The special role in the octet channel plays the mixing. There are two mixing models on the market now: the octet-singlet and quark-flavor mixing schemes. We reformulate these models without introducing the nonphysical states with indefinite masses. Each scheme implies a certain constraint for the meson decay constants (31),(32). Both mixing models are compatible only in the exact $SU(3)_f$ limit.

Using the data on the transition form factors of the η, η' mesons, the ASR allows to extract the set of decay constants in the octet-single, quark-flavor schemes as well as in the mixing-scheme independent way, if we add an additional constraint of ratio of radiation decays of J/ψ meson (41). It is shown, that the current data precision permits both, the octet-singlet and quark-flavor mixing schemes. The future improvement of experimental data on transition form factors of η, η' mesons and ratio $R_{J/\psi}$, expected from the BELLE and BES-III collaborations, can determine which scheme is more suitable for description of the mixing in the $\eta - \eta'$ system.

Acknowledgments

We thank $\boxed{\textit{A. Bakulev}}$, S. Brodsky, S. Eidelman, B. Ioffe, F. Jegerlehner, A. Kataev, D. Melikhov, S. Mikhailov, A. Pimikov, M. Polyakov, N. Stefanis and S. Uehara for discussions and useful comments. Y.K. thanks Y. Okada for a warm hospitality at KEK, where a part of this work was completed. A.O. and O.T. thank N. Stefanis and M. Polyakov for a warm hospitality during their stay at ITP, Ruhr-University Bochum. This work is supported in part by RFBR, research projects 12-02-00613a, 12-02-00284a, and by the Heisenberg-Landau program.

---

[1] J. S. Bell, R. Jackiw, Nuovo Cim. **A60**, 47-61 (1969).
[2] S. L. Adler, Phys. Rev. **177**, 2426-2438 (1969).
[3] A. D. Dolgov, V. I. Zakharov, Nucl. Phys. **B27**, 525-540 (1971).
[4] J. Horejsi, Phys. Rev. **D32**, 1029 (1985).
[5] J. Horejsi, O. Teryaev, Z. Phys. **C65**, 691-696 (1995).
[6] O. L. Veretin, O. V. Teryaev, Phys. Atom. Nucl. **58**, 2150-2155 (1995).
[7] Y. N. Klopot, A. G. Oganesian and O. V. Teryaev, Phys. Lett. B **695**, 130 (2011) [arXiv:1009.1120 [hep-ph]].
[8] Y. N. Klopot, A. G. Oganesian and O. V. Teryaev, Phys. Rev. D **84**, 051901 (2011) [arXiv:1106.3855 [hep-ph]]. Y. Klopot, A. Oganesian and O. Teryaev, Nucl. Phys. Proc. Suppl. **219-220** (2011) 141.
[9] Y. Klopot, A. Oganesian and O. Teryaev, JETP Lett. **94**, 729 (2011) [JETP Lett. **94**, 1 (2012)] [arXiv:1110.0474 [hep-ph]].
[10] D. Melikhov and B. Stech, Phys. Rev. D **85**, 051901 (2012) [arXiv:1202.4471 [hep-ph]].
[11] D. Melikhov and B. Stech, arXiv:1206.5764 [hep-ph].
[12] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **80**, 052002 (2009) [arXiv:0905.4778 [hep-ex]].
[13] P. del Amo Sanchez *et al.* [BABAR Collaboration], Phys. Rev. D **84**, 052001 (2011) [arXiv:1101.1142 [hep-ex]].
[14] S. Uehara *et al.* [Belle Collaboration], arXiv:1205.3249 [hep-ex].
[15] G. P. Lepage and S. J. Brodsky, Phys. Rev. **D22**, 2157 (1980).
[16] A. Khodjamirian, Int. J. Mod. Phys. A **25**, 513 (2010) [arXiv:0909.2154 [hep-ph]].
[17] S. V. Mikhailov and N. G. Stefanis, Mod. Phys. Lett. A **24**, 2858 (2009) [arXiv:0910.3498 [hep-ph]].
[18] A. P. Bakulev, S. V. Mikhailov, A. V. Pimikov and N. G. Stefanis, Phys. Rev. D **84**, 034014 (2011) [arXiv:1105.2753 [hep-ph]]. A. P. Bakulev, S. V. Mikhailov, A. V. Pimikov and N. G. Stefanis, Phys. Rev. D **86**, 031501 (2012) [arXiv:1205.3770 [hep-ph]]. A. V. Pimikov, A. P. Bakulev, S. V. Mikhailov and N. G. Stefanis, arXiv:1208.4754 [hep-ph].
[19] S. S. Agaev, V. M. Braun, N. Offen and F. A. Porkert, Phys. Rev. D **83**, 054020 (2011) [arXiv:1012.4671 [hep-ph]]. S. S. Agaev, V. M. Braun, N. Offen and F. A. Porkert, arXiv:1206.3968 [hep-ph].
[20] P. Kroll, Eur. Phys. J. C **71**, 1623 (2011) [arXiv:1012.3542 [hep-ph]].
[21] A. V. Radyushkin, Phys. Rev. D **80**, 094009 (2009) [arXiv:0906.0323 [hep-ph]].
[22] M. V. Polyakov, JETP Lett. **90**, 228 (2009) [arXiv:0906.0538 [hep-ph]].
[23] X. -G. Wu and T. Huang, Phys. Rev. D **82**, 034024 (2010) [arXiv:1005.3359 [hep-ph]]. X. -G. Wu and T. Huang, Phys. Rev. D **84**, 074011 (2011) [arXiv:1106.4365 [hep-ph]]. X. -G. Wu, T. Huang and T. Zhong, arXiv:1206.0466 [hep-ph].
[24] S. J. Brodsky, F. -G. Cao and G. F. de Teramond, Phys. Rev. D **84**, 033001 (2011) [arXiv:1104.3364 [hep-ph]]. S. J. Brodsky, F. -G. Cao and G. F. de Teramond, Phys. Rev. D **84**, 075012 (2011) [arXiv:1105.3999 [hep-ph]]. G. F. de Teramond and S. J. Brodsky, arXiv:1203.4025 [hep-ph].
[25] A. Stoffers and I. Zahed, Phys. Rev. C **84**, 025202 (2011) [arXiv:1104.2081 [hep-ph]].
[26] A. E. Dorokhov, Phys. Part. Nucl. Lett. **7**, 229 (2010) [arXiv:0905.4577 [hep-ph]]. A. E. Dorokhov, arXiv:1003.4693 [hep-ph].
[27] P. Kotko and M. Praszalowicz, Phys. Rev. D **81**, 034019 (2010) [arXiv:0912.0029 [hep-ph]].
[28] T. N. Pham and X. Y. Pham, Int. J. Mod. Phys. A **26**, 4125 (2011) [arXiv:1101.3177 [hep-ph]].
[29] D. McKeen, M. Pospelov and J. M. Roney, Phys. Rev. D

85, 053002 (2012) [arXiv:1112.2207 [hep-ph]].

[30] C. -C. Lih and C. -Q. Geng, Phys. Rev. C **85**, 018201 (2012) [arXiv:1201.2220 [hep-ph]]. C. -Q. Geng and C. -C. Lih, Phys. Rev. C **86**, 038201 (2012) [arXiv:1209.0174 [hep-ph]].

[31] P. Lichard, Phys. Rev. D **83**, 037503 (2011) [arXiv:1012.5634 [hep-ph]].

[32] E. Ruiz Arriola and W. Broniowski, Phys. Rev. D **81**, 094021 (2010) [arXiv:1004.0837 [hep-ph]].

[33] N. I. Kochelev and V. Vento, Phys. Rev. D **81**, 034009 (2010) [arXiv:0912.2172 [hep-ph]].

[34] Y. .M. Bystritskiy, V. V. Bytev, E. A. Kuraev and A. N. Ilyichev, Phys. Part. Nucl. Lett. **8**, 73 (2011) [arXiv:0912.3668 [hep-ph]].

[35] S. Noguera and V. Vento, Eur. Phys. J. A **46**, 197 (2010) [arXiv:1001.3075 [hep-ph]]. S. Noguera and S. Scopetta, Phys. Rev. D **85**, 054004 (2012) [arXiv:1110.6402 [hep-ph]].

[36] H. L. L. Roberts, C. D. Roberts, A. Bashir, L. X. Gutierrez-Guerrero and P. C. Tandy, Phys. Rev. C **82**, 065202 (2010) [arXiv:1009.0067 [nucl-th]].

[37] Z. -k. Guo and Q. Zhao, Eur. Phys. J. C **72**, 1964 (2012) [arXiv:1108.0241 [hep-ph]].

[38] W. Lucha and D. Melikhov, J. Phys. G **39**, 045003 (2012) [arXiv:1110.2080 [hep-ph]]. I. Balakireva, W. Lucha and D. Melikhov, Phys. Rev. D **85**, 036006 (2012) [arXiv:1110.6904 [hep-ph]]. I. Balakireva, W. Lucha and D. Melikhov, arXiv:1208.1025 [hep-ph].

[39] H. Czyz, S. Ivashyn, A. Korchin and O. Shekhovtsova, Phys. Rev. D **85**, 094010 (2012) [arXiv:1202.1171 [hep-ph]].

[40] S. Noguera and V. Vento, arXiv:1205.4598 [hep-ph].

[41] S. Dubnicka, A. Z. Dubnickova and A. Liptaj, Prog. Part. Nucl. Phys. **67**, 418 (2012).

[42] P. Masjuan, arXiv:1206.2549 [hep-ph].

[43] L. Rosenberg, Phys. Rev. **129**, 2786-2788 (1963).

[44] V. L. Eletsky, B. L. Ioffe and Y. I. Kogan, Phys. Lett. B **122**, 423 (1983).

[45] A. V. Radyushkin and R. T. Ruskov, Nucl. Phys. B **481**, 625 (1996) [arXiv:hep-ph/9603408].

[46] S. L. Adler, W. A. Bardeen, Phys. Rev. **182**, 1517-1536 (1969).

[47] G. 't Hooft in "Recent developments in gauge theories" ed. by G. 't Hooft et al., Plenum Press, New York NY (1980).

[48] F. Jegerlehner, O. V. Tarasov, Phys. Lett. **B639**, 299-306 (2006). [hep-ph/0510308].

[49] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **147**, 448 (1979).

[50] S. J. Brodsky, G. P. Lepage, Phys. Rev. **D24**, 1808 (1981).

[51] A. V. Radyushkin, Acta Phys. Polon. **B26**, 2067-2096 (1995). [hep-ph/9511272].

[52] H. J. Behrend *et al.* [CELLO Collaboration], Z. Phys. C **49**, 401 (1991).

[53] J. Gronberg *et al.* [CLEO Collaboration], Phys. Rev. D **57**, 33 (1998) [hep-ex/9707031].

[54] E. J. Schreier, Phys. Rev. D **3**, 980 (1971).

[55] G. T. Gabadadze and A. L. Kataev, JETP Lett. **61**, 448 (1995) [Pisma Zh. Eksp. Teor. Fiz. **61**, 439 (1995)] [hep-ph/9502384].

[56] J. Mondejar and K. Melnikov, arXiv:1210.0812 [hep-ph].

[57] K. G. Chetyrkin, S. Narison and V. I. Zakharov, Nucl. Phys. B **550**, 353 (1999) [hep-ph/9811275]. M. N. Chernodub, F. V. Gubarev, M. I. Polikarpov and V. I. Zakharov, Phys. Lett. B **475**, 303 (2000) [hep-ph/0003006].

[58] V. V. Anisovich, D. I. Melikhov, V. A. Nikonov, Phys. Rev. **D55**, 2918-2930 (1997). [hep-ph/9607215].

[59] T. Feldmann, P. Kroll, Eur. Phys. J. **C5**, 327-335 (1998). [hep-ph/9711231].

[60] T. Feldmann, P. Kroll and B. Stech, Phys. Rev. D **58**, 114006 (1998) [hep-ph/9802409]. Phys. Lett. B **449**, 339 (1999) [hep-ph/9812269].

[61] T. Feldmann, Int. J. Mod. Phys. A **15**, 159 (2000) [hep-ph/9907491].

[62] H. Leutwyler, Nucl. Phys. Proc. Suppl. **64**, 223 (1998) [hep-ph/9709408]. R. Kaiser and H. Leutwyler, In *Adelaide 1998, Nonperturbative methods in quantum field theory* 15-29 [hep-ph/9806336].

[63] P. Herrera-Siklody, J. I. Latorre, P. Pascual and J. Taron, Nucl. Phys. B **497**, 345 (1997) [hep-ph/9610549].

[64] J. Schechter, A. Subbaraman and H. Weigel, Phys. Rev. D **48**, 339 (1993) [hep-ph/9211239].

[65] J. Beringer *et al.* [Particle Data Group Collaboration], Phys. Rev. D **86**, 010001 (2012).

[66] V. A. Novikov, M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **165**, 55 (1980).

[67] P. Ball, J. M. Frere, M. Tytgat, Phys. Lett. **B365**, 367-376 (1996). [hep-ph/9508359].

[68] F. -G. Cao and A. I. Signal, Phys. Rev. D **60**, 114012 (1999) [hep-ph/9908481].

[69] R. Escribano, J. -M. Frere, JHEP **0506**, 029 (2005). [hep-ph/0501072].

[70] T. Feldmann, P. Kroll, Phys. Rev. **D58**, 057501 (1998). [hep-ph/9805294].

[71] In case of singlet channel ($a = 0$) this property is violated because of gluonic anomaly

[72] The growth of the octet combination is due to the fact, that $Q^2 F_{\eta\gamma}$ at $Q^2 > 20GeV^2$ slightly tends to go up, and $Q^2 F_{\eta\gamma}$ tends to go down, while f_η^8 is positive and f_η^8 is negative.

[73] Note, that in our approach the same relation exists only for the fields (see Eq. (26)).